



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Kobbernagel and colleagues have demonstrated that the course of PCD lung disease can be improved using a widely available and relatively low-cost pharmacological treatment. Indeed, lung function remained unchanged over the intervention period, and a treatment beyond 6 months is needed to evaluate the efficacy in this field. In one study, FEV₁ decline among several patients with cystic fibrosis in a French Registry appeared less marked after than before prolonged azithromycin treatment initiation.⁹ However, in that non-randomised, retrospective study the decision to start long-term azithromycin was based on a pre-treatment FEV₁ level that was quite low, thus more sick patients with cystic fibrosis were included. This factor is a striking difference with the BESTCILIA trial that ruled out PCD patients with more severe pulmonary disease.

Kobbernagel and colleagues are to be congratulated for this well designed study including a large number of patients, which is difficult to achieve in rare disorders. The participating centres are members of either the European Reference Network on respiratory diseases or the Better Experimental Approaches to Treat PCD network.^{10,11} The role of networks for uncommon diseases is crucial to improve access of patients to diagnosis and treatment, and to encourage collaborative research from different countries.

Future directions for research on treatment of PCD airways infections should include longer prospective clinical trials, the recruitment of patients with more impaired lung function and those with *Pseudomonas* spp colonisation. Comparative studies with prophylactic nebulized or intermittent systemic

antibiotics are also necessary. The PCD community is confident that BESTCILIA is only the first of a series of future multicentre clinical trials that will provide more guidance for precision medicine for the PCD population.

FS is member of Better Experimental Approaches to Treat PCD (COST Action 1407), and the affiliated Department is part of the European Reference Network on respiratory diseases. All other authors declare no competing interests.

Marco Poeta, Melissa Borrelli, *Francesca Santamaria
santamar@unina.it

Department of Translational Medical Sciences, Section of Pediatrics, Federico II University, 80131 Naples, Italy

- 1 Mirra V, Werner C, Santamaria F. Primary ciliary dyskinesia: an update on clinical aspects, genetics, diagnosis, and future treatment strategies. *Front Pediatr* 2017; **5**: 135.
- 2 Shoemark A, Dell S, Shapiro A, Lucas JS. ERS and ATS diagnostic guidelines for primary ciliary dyskinesia: similarities and differences in approach to diagnosis. *Eur Respir J* 2019; **54**: 1901066.
- 3 Poeta M, Maglione M, Borrelli M, Santamaria F. Non-cystic fibrosis bronchiectasis in children and adolescents: Neglected and emerging issues. *Pediatr Neonatol* 2019; published online Oct 11. DOI:10.1016/j.pedneo.2019.09.013.
- 4 Kobbernagel HE, Buchvald FF, Haarman EG, et al. Efficacy and safety of azithromycin maintenance therapy in primary ciliary dyskinesia (BESTCILIA): a multicentre, double-blind, randomised, placebo-controlled phase 3 trial. *Lancet Respir Med* 2020; **8**: 493–505.
- 5 Kudoh S, Azuma A, Yamamoto M, Izumi T, Ando M. Improvement of survival in patients with diffuse panbronchiolitis treated with low-dose erythromycin. *Am J Respir Crit Care Med* 1998; **157**: 1829–32.
- 6 Bush A. Azithromycin is the answer in paediatric respiratory medicine, but what was the question. *Paediatr Respir Rev* 2019; published online Aug 16. DOI:10.1016/j.prrv.2019.07.002.
- 7 Lucas JS, Davis SD, Omran H, Shoemark A. Primary ciliary dyskinesia in the genomics age. *Lancet Respir Med* 2020; **8**: 202–16.
- 8 Haendel MA, Chute CG, Robinson PN. Classification, ontology, and precision medicine. *N Engl J Med* 2018; **379**: 1452–62.
- 9 Denis A, Touzet S, Diabaté L, et al. Quantifying long-term changes in lung function and exacerbations after initiation of azithromycin in cystic fibrosis. *Ann Am Thorac Soc* 2020; **17**: 195–201.
- 10 Humbert M, Wagner TO. Rare respiratory diseases are ready for primetime: from Rare Disease Day to the European Reference Networks. *Eur Respir J* 2017; **49**: 1700085.
- 11 Rubbo B, Behan L, Dehlink E, et al. Proceedings of the COST action BM1407 inaugural conference BEAT-PCD: translational research in primary ciliary dyskinesia - bench, bedside, and population perspectives. *BMC Proc* 2016; **10**: 66.



Understanding pathways to death in patients with COVID-19

Published Online
April 6, 2020
[https://doi.org/10.1016/S2213-2600\(20\)30165-X](https://doi.org/10.1016/S2213-2600(20)30165-X)
See Review page S06

Since the first cases of coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), were identified in China in December, 2019, we have witnessed increasing numbers of infections and associated deaths worldwide. Although the case fatality rate for SARS-CoV-2 infection (ie, the total number of deaths in patients positive for SARS-CoV-2 divided by the total

number of people with a positive test) is not high, given the huge scale of the pandemic, the actual numbers of deaths are considerable.

In *The Lancet Respiratory Medicine*, Jason Phua and colleagues¹ provide an excellent overview of the current issues raised by COVID-19—in particular, the impact of the disease on intensive care. The Review is clearly and comprehensively written, covering many aspects of the




Figure: Possible paths to death and recovery in patients needing respiratory support

CPAP=continuous positive airway pressure. ECMO=extracorporeal membrane oxygenation. NIV=non-invasive ventilation.

disease, from epidemiology and diagnosis through to intensive care treatment and resource management. One issue raised by this Review is how the reported case fatality rates for patients with COVID-19 can be accurately interpreted.

Currently reported case fatality rates vary from 1% to more than 7%,² but these values must be interpreted with caution. For example, where massive screening has been performed in the whole population (eg, in South Korea and Switzerland), overall case fatality rates of less than 1% have been reported, because the denominator included many mild or asymptomatic cases. However, in countries where only people requiring hospital admission are being screened (eg, Italy and Spain), case fatality rates have exceeded 5%, because the denominator is much smaller.

The actual cause of death is also important in interpreting case fatality rates. Respiratory failure is obviously the main cause,³ as was also the case in previous viral pandemics, such as the Spanish flu of 1918. Today, however, many patients can be supported by invasive mechanical ventilation until the lungs recover. If the situation deteriorates, use of extracorporeal membrane oxygenation (ECMO) systems can control gas exchange for weeks. COVID-19 is sometimes complicated by shock and multiple organ failure,^{4,5} but the real course of the disease is not yet well described. Knowing that non-survivors are more likely to have low lymphocyte counts or high C-reactive protein or D-dimer levels^{3,6,7} provides no information about the actual process of death. The precise role of secondary bacterial infections has also not been well defined.

Ethical issues also have a relevant role in interpreting case fatality rates, especially when the elderly and frail are more at risk and when resources are stretched so that some

Panel: Three possible characteristics of the dying process in COVID-19

Predominant terminal organ failure

- Terminal respiratory failure: mechanical ventilation and ECMO used
- Terminal respiratory failure: mechanical ventilation used, ECMO available but not used
- Terminal respiratory failure: mechanical ventilation used, ECMO not available
- Respiratory failure: mechanical ventilation available but not used
- Respiratory failure: mechanical ventilation hardly or not available
- Septic shock, multiple organ failure
- Cardiogenic shock (acute myocardial injury or myocarditis)
- Other

Proportionality of care in the dying process

- Withholding life support: life support available but considered to be disproportionate; life support hardly available (significant constraints)
- Withdrawing life support
- Full care but no cardiopulmonary resuscitation
- Full care including cardiopulmonary resuscitation

Involvement of COVID-19 in the dying process

- Death attributed only to COVID-19 (previously healthy, predicted long life expectancy)
- Death primarily due to old age, frailty, or advanced disease (COVID-19 is an epiphenomenon)
- Death due to COVID-19 in an individual with limited life expectancy

COVID-19=coronavirus disease 2019. ECMO=extracorporeal membrane oxygenation.

form of rationing or triage might become necessary. In such a scenario, differentiating whether the cause of death is specifically due to COVID-19 or the result of treatment limitations can be difficult. Among patients who die before reaching the hospital, some will present too late in the course of the disease to be saved, whereas end-of-life care will be viewed as preferable for others because little chance of survival with a meaningful quality of life exists. In some patients, this decision might be influenced by known individual preferences. Similarly, not all critically ill patients in hospital will be admitted to the intensive care unit (ICU), because the chances of meaningful survival for some will be viewed as too low;⁸ for these patients, non-invasive

ventilation and perhaps even some vasopressor support could be provided in so-called middle care units, but in case of further deterioration, mechanical ventilation will not be considered and death will occur because of severe hypoxaemia. As noted by Phua and colleagues,¹ a quarter of patients who died early in the Wuhan, China, outbreak did not receive invasive ventilation.⁹ Patients who deteriorate despite mechanical ventilation can be placed on resource-intensive ECMO systems (figure).¹⁰ The decision not to use ECMO might be made because the support system is not available or because such care is considered to be disproportionate in the context of limited staff numbers. The same considerations might apply to patients who develop renal failure. Use of renal replacement therapies is uncommon in those with COVID-19,^{4,6,7} although acute kidney injury might occur in a third of patients.

Several different scenarios can thus be considered when interpreting deaths from COVID-19. First, for patients admitted to the ICU, death might occur despite full intensive care support, including mechanical ventilation, ECMO, vasopressors, and renal replacement therapy. On modern ICUs, such deaths are expected to be infrequent; however, robust estimates of the number of deaths cannot be made from the mostly descriptive reports currently available. A second possible scenario for ICU and hospital patients is related to limitation of life-sustaining therapies because of poor predicted outcomes associated with old age, frailty, comorbidities, or profound disability, or because of effects of distributive limitations associated with lack of personnel, beds, or materials. A combination of these two factors often exists. A third scenario relates to patients admitted to the ICU or hospital whose deaths are not directly related to COVID-19. Especially in areas with high infection rates, patients might be admitted to the ICU with, for example, severe trauma or acute brain injury, test positive for SARS-CoV-2 during the ICU stay, and eventually die because of the initial injury; these deaths will still be attributed to COVID-19 and included in the statistics. Similarly, some patients might have SARS-CoV-2 infection, but the actual contribution of the virus to the patient's death might be minimal. For example, in a patient with metastatic cancer or terminal organ failure, is the viral infection or the patient's underlying condition the cause of death? The actual role of SARS-CoV-2 infection in such deaths is particularly difficult to evaluate in countries where only one cause can be reported on a death certificate.

To be able to better interpret case fatality rates, more data are thus needed (panel). First, the type and severity of organ failure: what are the real contributions of respiratory or cardiovascular failure? How many patients died with isolated respiratory failure, in shock or with acute kidney injury or multiple organ failure? Second, the actual process of death, including therapeutic limitation when present and the relative contributions of patient factors (eg, age and comorbidities) or environmental factors (eg, lack of facilities, beds, personnel, or equipment). Last, the real contribution to death of SARS-CoV-2 infection, because COVID-19 can be an epiphenomenon in some patients.

We have learnt a lot in a relatively short period of time, and the Review by Phua and colleagues¹ summarises this knowledge well. However, we still have a lot to learn. Among the many unanswered questions is the key issue related to the actual process leading to death. Global numbers of deaths and case fatality rates provide only crude information.

We declare no competing interests.

*Jean-Louis Vincent, Fabio S Taccone
jlvincen@intensive.org

Department of Intensive Care, Erasme University Hospital, Université Libre de Bruxelles, 1070 Brussels, Belgium

- 1 Phua J, Weng L, Ling L, et al. Intensive care management of coronavirus disease 2019 (COVID-19): challenges and recommendations. *Lancet Respir Med* 2020; published online April 6. [https://doi.org/10.1016/S2213-2600\(20\)30161-2](https://doi.org/10.1016/S2213-2600(20)30161-2).
- 2 Onder G, Rezza G, Brusaferro S. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA* 2020; published online March 23. DOI:10.1001/jama.2020.4683.
- 3 Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* 2020; published online March 3. DOI:10.1007/s00134-020-05991-x.
- 4 Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; **395**: 507–13.
- 5 Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020; **323**: 1061–69.
- 6 Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; **395**: 1054–62.
- 7 Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020; published online Feb 28. DOI:10.1056/NEJMoa2002032.
- 8 Emanuel EJ, Persad G, Upshur R, et al. Fair allocation of scarce medical resources in the time of Covid-19. *N Engl J Med* 2020; published online March 23. DOI:10.1056/NEJMs2005114.
- 9 Xie J, Tong Z, Guan X, Du B, Qiu H, Slutsky AS. Critical care crisis and some recommendations during the COVID-19 epidemic in China. *Intensive Care Med* 2020; published online March 2. DOI:10.1007/s00134-020-05979-7.
- 10 Ronco C, Navalesi P, Vincent JL. Coronavirus epidemic: preparing for extracorporeal organ support in intensive care. *Lancet Respir Med* 2020; **8**: 240–41.